

Regular quantum interiors for black holes

Emilio Elizalde^{*,†} and Sergi R. Hildebrandt

Instituto de Ciencias del Espacio (CSIC)

È Institut d'Estudis Espacials de Catalunya (IEEC/CSIC)

Edifici Nexus, Gran Capità 2-4, 08034 Barcelona, Spain

A family of spacetimes suitable for describing the interior of a non-rotating black hole is constructed. The stress-energy tensor is that of a spherically symmetric vacuum, as commonly assumed nowadays. The problem of matching the exterior with the interior region is solved exactly, without using any massive shell nor having to restrict oneself to only asymptotically well-behaved solutions, whatsoever. The main physical and geometrical properties of the resulting black hole solutions are described. As models for the interior the general solution found includes, in particular, two known previous attempts at solving the problem. Finally, effective macroscopic properties of the solution are linked with quantization issues of the corresponding spacetime.

Work on collapsing bodies has shown [1,2] that there exists a mass above which the collapse of an object and the appearance of a singularity are unavoidable. In searching for new solutions, with the aim to circumvent such conclusion, it has been claimed that other quantum effects, specially those associated with the quantum vacuum, might actually yield regular solutions everywhere inside the object. Indeed, to stop gravitational collapse of a very massive body, the final solution should develop a static region that coincides with the core of the object. The strong energy conditions [3] must be violated inside the final body, in order to make the singularity avoidable. Even though any realistic object may differ significantly from spherical symmetry at the beginning of the collapse, there are several mechanisms by which, close to the final state, spherical symmetry holds, both for the exterior as well as for the interior regions [4,5], to the point where the classical notions of space and time lose their meaning.

The general expression for the matter-energy content of such contributions, for spherically symmetric bodies, has the form $\rho + p = 0$, and $p_2 = p_3$ [6,7], where ρ is the mass-energy density, p the radial stress (or pressure) and $p_2 = p_3$ the transversal stresses (pressures) measured by any local observer (proper reference frame).

We will here deal with the family of maximal spherically symmetric spacetimes expanded from flat spacetime by a geodesic radial null one-form (GRNSS spaces, see [8–10]), a subfamily of Kerr-Schild (KS) metrics. They are given by

$$ds^2 = ds_\eta^2 + 2H(r)\ell \otimes \ell, \quad (1)$$

where ds_η^2 stands for the flat spacetime metric, H is an arbitrary function of r , the radial coordinate of the spherical symmetry —defined in some open region of the manifold— and ℓ is a geodesic radial null one-form.

These metrics include *all* spherically symmetric spacetimes which have a static region and satisfy the previous energy-matter constraints. Another expression for the family is (with signature $(-+++)$)

$$ds^2 = -(1-H)dt^2 + 2Hdt\,dr + (1+H)dr^2 + r^2(d\theta^2 + \sin^2\theta\,d\varphi^2), \quad (2)$$

with $\ell = (1/\sqrt{2})(dt + dr)$. The other possibility, i.e. $\ell = (1/\sqrt{2})(dt - dr)$ yields the same physical results. Since $H = H(r)$, ∂_t is an integrable Killing vector.

The above choice avoids coordinate problems near their possible horizons, e.g. when $H = 1$.

In the region where $H < 1$, the existence of such integrable Killing vector allows to write the whole family of metrics in an explicitly static form,

$$ds^2 = -(1-H)dt_s^2 + \frac{1}{1-H}dr^2 + r^2(d\theta^2 + \sin^2\theta\,d\varphi^2),$$

where dt_s is related to dt by $dt_s = dt - [H/(1-H)]dr$. This last expression for the family looks as a generalization of the well known Schwarzschild metric and may help to identify the class of spacetimes we are dealing with, [11].

In order to recover the natural scheme of collapsed objects, one must consider the matching of two spherically symmetric solutions of the Einstein equations. Here, the interior solution will belong to the class GRNSS while the exterior one may be *any* solution for a non-rotating classical black hole. One can consider e.g. a charged, Reissner-Nordström solution, a Schwarzschild solution [2,12,13], or even a cosmological background including a cosmological constant, [14]. Regardless of the choice of exterior metric, we have the additional advantage that *all* these exterior solutions also belong to the GRNSS family of spaces. As a consequence, the matching becomes simple.

There are in fact many possible choices for the equations defining the matching hypersurface [9], the most interesting is simply to choose the hypersurface of constant r . Other possibilities (avoiding the singularity) should converge to this choice.

The corresponding matching conditions translate into: $[H] = 0$, $[H'] = 0$, where $[f] = f_{\text{ext}} - f_{\text{int}}$ and the prime stands for the radial ordinary derivative. Their physical interpretation is direct, the first is the continuity of the mass function, while the second assures the continuity of radial pressures.

To avoid the central singularity we could impose a specific form of the matter-energy content under some general accepted scheme of quantum behaviour close to the

origin. However it is more interesting to demand only that the solution be regular at the origin and study the main (universal) properties of all such candidates. In this way we shall extract some general information, unbiased by our prejudices on the “ultimate state of matter”.

A first remarkable consequence is that, regardless of the chosen specific model for the interior, *all* become isotropic near the origin, i.e. $p_2 = p$ at the origin. This property has been always imposed ever since the incorporation of high energy quantum effects in the field, and is tacitly assumed in all previous attempts. Here it *directly follows* from the GRNSS defining conditions, for which we only demanded $\rho = -p$, $p_2 = p_3$, and partial staticity, but not necessarily isotropy. In consequence, we do get a *smooth* transition from the exterior phase to the central one, as expected by previous authors, see e.g. the conclusions in [16]. Moreover -and this is also a very important point- the need for a (thin) layer has now disappeared, [8]. In summary, the matching of two GRNSS spaces recovers, in an absolutely natural way, *all* the features of existing collapse schemes for massive objects and extends this issue to the domain of new highly energetic quantum processes.

For a solution of the GRNSS class to be everywhere regular, it suffices that $H_2(r=0) = 0$, and $H'_2(r=0) = 0$, where label 2 refers to the interior solution. This result on the regularity of the inner solution is based on the analysis of the Riemannian invariants of the metrics. For instance, $\mathcal{R}^2 \equiv R_{\mu\nu\lambda\rho}R^{\mu\nu\lambda\rho}$ yields $(H'')^2 + 4(H'/r)^2 + 4(H/r^2)^2$. Any regular interior model develops here a maximum limiting value for all these invariants, as also expected by several authors, e.g. [18].

Another important consequence of the general regular models is that the “mass function”, defined as usual by $m(r) = r(1 - g_{static}^{rr})/2$, is $rH(r)/2$ and becomes smaller and smaller as the origin is approached, until it vanishes. Thus *all* these models also yield the result that they are asymptotically free in the sense that the gravitational charge $m(r)$ vanishes at the origin. In [19], Sect. II, results were claimed to be possibly valid also for any asymptotically free model of the interior core. Here we prove rigorously that *all* regular-at-the-origin GRNSS spaces are asymptotically free and share those desirable physical properties.

A number of papers have dealt with this issue, looking for plausible energy contributions near the origin. Thus, in [20] it is advocated for a de Sitter core, which has been hitherto the most studied situation, $H = \Lambda r^2/3$, clearly satisfying the regularity conditions. But there is still a broad open window for other alternatives. As yet there is no quantum gravity theory available and therefore an assumption regarding the precise form of the stress-energy tensor at the origin cannot be made. However there are impelling arguments in favor of such a behaviour near the origin and the de Sitter choice will then correspond to the limiting case.

For the exterior region, one usually assumes a Schwarzschild black hole solution, i.e. a non-charged one (the

case of a charged black hole can be easily recovered from our results below). Observational evidence [21], as well as theoretical arguments [12], lead, on the other hand, to the conclusion that non-charged black holes are the most common ones in our universe. Nevertheless, one can also consider the effect of adding a cosmological background to the Schwarzschild solution in terms of a cosmological term, and/or also other contributions, as for instance quantum contributions coming from the vacuum polarization of the exterior region close to the matching hypersurface. Generically one will have a relation of the type $H_2(r \sim 0) \sim \Lambda_2 r^2/3$, and $H_1 = H_1(r, m_1, \Lambda_1, \{\alpha_i^1\})$, where m_1 is the gravitational mass as measured by an external observer, Λ_1 the (possible) cosmological term of the outer region, and $\{\alpha_i^1\}$ a set of parameters describing the strength of other effects, such as the charge of the black hole, vacuum polarization of the exterior solution, etc. By virtue of the matching conditions we will have schematically $H_2 = H_2(r; m_1, \Lambda_1, \{\alpha_i^{1,2}\}, R)$, where R is the value at which the two solutions match. The energy-matter density for any metric of the GRNSS spaces reads $(G=c=1)8\pi\rho_2 = (H_2 r')/r^2$. Thus ρ_2 will have an analogous dependence which, once imposed the criteria $8\pi\rho_2(r=0) = \Lambda_2$, will yield a relationship of the type $\Lambda_2 = F(R, m_1, \Lambda_1, \{\alpha_i^{1,2}\})$. In order to determine R in terms of the rest of the parameters, one has to consider a specific model for the interior. As in [1,2,12], we encounter a big set of allowed models for the interiors of the objects given a common exterior and the ultimate task is that of finding realistic physical models for these interiors. In [9] we perform in detail the calculations of the corresponding models in our formulation, as well as of a whole set of new proposals, in order to investigate the universality of the conclusions derived from these particular examples.

The first known model [4,19] has its analogue here taking $H_1 = 2m_1/r - (1/3) \times (\alpha m_1/r^2)^2$, where α^2 is a number related with the number and types of the quantized fields, being of order unity in Planckian units, and $H_2 = r^2/3B(B + Cr^3)$, where $B = \alpha/(6 - \alpha^2 m_1/R^3)$ and $C = (2/\alpha m_1)[1 - 3/(6 - \alpha^2 m_1/R^3)]$. H_1 takes into account corrections coming from the polarization of the vacuum that is expected to be proportional to $\mathcal{R}^2$. The second known model [23,7] is retrieved by setting $H_1 = 2m_1/r + \Lambda_1 r^2/3$, and $H_2 = (R^2/3)\{(\beta/\alpha)[\exp(-\alpha\tilde{r}^3) - 1]/\tilde{r} - \gamma\tilde{r}^2\}$, where $\gamma = -\beta \exp(-\alpha) - \Lambda_1$, and α, β are solutions of transcendental equations. A valid approximate expression is $\alpha \simeq (\Lambda_1 - \Lambda_2)R^3/6m_1$, and $\beta \simeq \Lambda_1 - \Lambda_2$. The other set of new models considered is described by $H_1 = 2m_1/r + \Lambda_1 r^2/3$, and $H_2 = b_2 r^2 + b_N r^N$, with $N \geq 3$, where b_2 is given by $(2m_1/R)(N+1)/(N-2) + \Lambda_1 R^2/3$, and $b_N = -6m_1/R(N-2)$.

Obviously the value of R will depend in general on the regularization scale at which the usual concepts of space and time lose their sense. To fix ideas, let us assume that the regularization scale is of Planckian order (other choices are to be found in [9]). The result is that in

all the preceding models R remains constrained to the range $2k\sqrt[3]{M}(10^{-20}\text{cm})$, with $k \sim [1, 10]$, and M defined by $m_1/m_\odot$, being $m_\odot$ the solar mass.

There are some issues worth mentioning about the models. First, all possess a matching R which is well above the regularization scale for any astrophysical object ($m \sim m_\odot$). For instance, in the case of the Planck scale, the result is $R/L_{Pl} \sim 10^{12} \gg 1$. This boosts our confidence on the plausibility of the solutions found, since they are indeed very far from the scales where the applied scheme of continuous space and time loses its sense. On the other hand, these objects exhibit in general two main horizons. The first is the usual event horizon, that coincides with the typical one of a black hole, located at $r = 2m_1$. The second is a (non-global) Cauchy horizon (CH) [3,2] near the origin, where $H_2 = 1$. For $r < r_{\text{Cauchy}}$ the spacetime becomes static and we enter into the de Sitter core. In addition to these “local” horizons, we could also have others, coming from the cosmological background [7,22,9] (see also Fig. 1 in [9]). The total number of horizons affects the so-called stability of the inner CH. This question has received a lot of attention in the literature, where non-stationary perturbations of the static, or stationary, models show, at the classical or semiclassical level, that the CH may become unstable, e.g. [5]. Yet this (dynamical) instability is weak [15], so that the formation of the core is still possible. Moreover, the strong energy conditions [3,13] are violated far enough from the regularization scale, so that eventually the singularity is not created. This is in accordance with known singularity theorems, which need the strong energy condition to be satisfied near the singularity.

Also worth considering is the stability of the solutions: will they still exist in case of changes in the parameters such as m_1 , because of mass accretion, the regularization scale, etc.? If our models were highly dependent on some of these parameters, then the regular solution would be valid only for a “fine tuned configuration”. Along the lines of [16] we have proven that our models are stable. Another issue concerns the creation of a closed world inside the collapsed object. This possibility can be interpreted as the creation of a new (inflationary) macroscopic universe connected with the collapsed body via its core (see Fig. 2 in [9]). In the two models studied in [19,7,22] this property holds. Here, because of the properties of asymptotically free cores and the existence of a non-global Cauchy horizon this possibility is always present, see also [17].

We note that all these models keep the matching hypersurface inside the event horizon. In general, this has to be indeed the case for any collapsed body exceeding approximately three solar masses. Of course, other objects with a lower mass can also form a black hole provided its matter-energy content is sufficiently compressed. Therefore, in order to analyze whether the singularity of a black hole is avoidable, we must descend to the physics inside the event horizon and study matter-energy contents that will allow for the necessary violation of the energy con-

ditions to stop the creation of the singularity. So far the only candidates are those coming from quantum field theory, because we fairly know that matter and energy become quantized under the physical conditions prevailing in these objects. Even though a theory of quantum gravity is yet unavailable and, therefore, no exact solution is still known for such merge, there are some results that point towards a plausibility of the avoidance of singularities. The reasons for such confidence are twofold. First, concerning the type of energy contributions to be expected near the origin of the object, several studies [20] lead to the conclusion that the quantum energy of the vacuum acts as a negative stress and could remove the appearance of the singularity. This crucial point is explicitly present in our geometrical description in terms of the isotropization near the origin and through its association with an internal Λ . These first results touch the most intricate problem, namely that of the plausible physics near the regularization scale, but let us recall that different such possibilities can be easily incorporated into our scheme, e.g. stringy black holes [9].

Second, there is the region far away from the renormalization scale, where one trusts the semiclassical theory of gravity [23] to be an accurate enough description of the involved physics. In fact this can be proven for any object satisfying $m_1 \gg M_{\text{Reg}}$, $R/L_{\text{Reg}} \gg 1$. For instance, for an astrophysical black hole and a regularization scale of the order of the Planck scale, we obtain $R/L_{Pl} \sim 10^{12}$, $R \sim 10^{-20}\text{cm}$, see also [4]. Thus, the fundamental question appears of whether any of these models corresponds to a solution of the semiclassical equations of gravity, at least in the domain where r/L_{Reg} is big enough. As far, no one has found any such suitable model, however under very general assumptions we have considered here *all* the possible candidates with spherical symmetry which have a static region and exhibit the expected behaviour for the quantum stress-energy tensor of a collapsed body. Thus, if any particular solution is found in the future, we believe that it will be inside this family.

To find the sources for the models here presented is not easy, but this will be a necessary step to undertake, to show the plausibility of avoidance of the singularities. To reach this goal within our scheme, one needs in fact to quantize the sources. The symmetries of the matter-energy tensor, i.e. $\rho + p = 0$, and the one coming from spherical symmetry, $p_2 = p_3$ are fundamental in order to check which type of quantum fields could correspond to their sources. Taking into account the spherical symmetry of the spacetime and, consequently, of the Einstein tensor, one easily checks that all classical fields adapted to the spherical symmetry satisfy this condition, see for instance Sect. 3.8 of [23]. This bonus was indeed expected, because spherical symmetry is a basic geometrical symmetry in the scheme. On the other hand, the other eq., $\rho + p = 0$, can be rewritten in a general covariant way, without referring to any special set of observers. Its expression is then $T_{ll} \equiv T_{\mu\nu}l^\mu l^\nu = 0$, where $\vec{l}$

is the geodesic radial null direction characteristic of the GRNSS spaces. In the scalar case, for instance, one obtains: $T_{ll} = (1 - 2\xi)\phi'^2 - 2\xi\phi\phi''$, where $(\cdot)' \equiv \vec{\ell}(\cdot)$, and ξ is a constant representing the coupling between the scalar and the gravitational field. If one imposes free scalar field equations, one finds no solution. The end result is that no free classical field satisfies this requirement. On the other hand, de Sitter spacetime has $\langle T_{\alpha\beta} \rangle = -\Lambda g_{\alpha\beta}$, thus $\langle T_{ll} \rangle = 0$, what means that the cancellation of T_{ll} is indeed due to quantum regularization of the classical field. A starting program of quantization of GRNSS spaces has led to the result that T_{ll} plays essentially the same role as T_{λ}^{λ} in related conformal spacetimes, and that the vanishing of $\langle T_{ll} \rangle$ is most likely an anomaly effect, also influenced by the mass of the field and its coupling to the gravitational field. Other sources are to be found within effective actions of M-theory or string theory, see e.g. [24].

Another —maybe less fundamental— way of attaining the expected quantum corrections would be to consider some anisotropic version of spherical collapse [6] and to perform the calculations associated with vacuum polarization effects. The form of the GRNSS spaces allows in fact for a direct calculation of such effects. This is a different physical point of view from the one advocated in [5], where the authors have been mainly concerned with the radiative processes inside the black hole. In our analysis, the results for the de Sitter and Schwarzschild cases yield well-known statements that have been duly taken into account in the models presented above.

We conclude that it seems natural that a suitable spacetime inside the GRNSS family will correspond to a physically realistic quantum model of the interior of a BH, free of singularities. On top of this conclusion, we recall again that other possibilities, even different singular models, are also contained in the GRNSS family —as long as they satisfy the usual stress-energy conditions for a spherical vacuum. In all, a remarkable amount of plausible realizations within this family. An extension to rotating BH in a similar framework has been recently considered [25].

Work supported by DGICYT (Spain), project PB96-0925 and CIRIT (Generalitat de Catalunya), contract 1999SGR-00257.

- bridge, 1973) .
- [4] E. Poisson and W. Israel, *Class. Quantum Grav.* **5**, L201 (1988).
 - [5] R. Balbinot and E. Poisson, *Phys. Rev. Lett.* **70**, 13 (1993); W. G. Anderson, P. R. Brady, W. Israel and S. M. Morsink, *Phys. Rev. Lett.* **70**, 1041 (1993). W. Israel, in *Black Holes and Relativistic Stars*, ed. R. M. Wald (The University of Chicago Press, Chicago, 1998).
 - [6] Ya. B. Zeldovich and I. D. Novikov, *The Structure and Evolution of the Universe* (Chicago University Press, Chicago, 1983).
 - [7] I. Dymnikova, *Gen. Rel. Grav.* **24**, 235 (1992)
 - [8] Magli, *G. Rept. Math. Phys.*, **44**, 407 (1999)
 - [9] E. Elizalde and S. R. Hildebrandt, (to be published).
 - [10] S. R. Hildebrandt, (to be published).
 - [11] R. C. Myers, and J. Z. Simon, *Phys. Rev. D* **38** 2434 (1988); J. T. Wheeler, *Nucl. phys.* **B273**, 732 (1986).
 - [12] C. W. Misner, K. S. Thorne and J. A. Wheeler, *Gravitation* (Freeman, San Francisco, 1973).
 - [13] R.M. Wald *General Relativity* (The University of Chicago Press, Chicago, 1984).
 - [14] D. Kramer, H. Stephani, M. MacCallum, and E. Herlt, *Exact Solutions of Einstein's Field Equations* (Cambridge Univ. Press, Cambridge, 1980) p. 156.
 - [15] A. Ori, *Phys. Rev. Lett.* **83**, 5423 (1999); *ibid* **68**, 2117 (1992); *ibid* **67**, 789 (1991). And references therein.
 - [16] R. Balbinot and E. Poisson, *Phys. Rev. D* **41**, 395 (1990).
 - [17] A. Borde, *Phys. Rev. D*, **55**, 7615 (1997).
 - [18] M. A. Markov, *Pis'ma Zh. Eksp. Teor. Fiz.* **36**, 214 (1982) [*Sov. Phys. JETP Lett.* **36**, 265 (1982)].
 - [19] V. P. Frolov, M. A. Markov and V. F. Mukhanov, *Phys. Rev. D* **41**, 383 (1990).
 - [20] E. B. Gliner, *Sov. Phys. JETP* **22**, 378 (1966); A. A. Starobinsky, *Phys. Lett.* **91B**, 99 (1980); M. A. Markov, *Ann. Phys., NY* **155**, 333 (1984).
 - [21] M. J. Rees, in *Black Holes and Relativistic Stars*, ed. R. M. Wald (The University of Chicago Press, Chicago, 1998).
 - [22] I. Dymnikova and B. Soltyssek, *Gen. Rel. Grav.* **30**, 1775 (1998).
 - [23] N. D. Birrel and P. C. W. Davies, *Quantum Theory in Curved Spaces* (Cambridge University Press, Cambridge, England, 1982).
 - [24] E. Ayón-Beato, and A. A. García, *Phys. Lett., B* **464**, 25 (1999), and references therein.
 - [25] Burinskii, A., [hep-th/0008129](#). Also in the Proceedings of the MG9, Rome (2000), to appear.

* elizalde@ieec.fcr.es, eli@ecm.ub.es

† <http://www.ieec.fcr.es/cosmo-www>

- [1] K. Schwarzschild, *xxx Ann. Phys.*, (1916); J.R. Oppenheimer and H. Snyder, *Phys. Rev.* **56**, 455 (1939).
- [2] S. Chandrasekhar, *The Mathematical Theory of Black Holes* (Clarendon Press, Oxford, 1983).
- [3] S. W. Hawking and G. F. R. Ellis, *The Large Scale Structure of Space-Time* (Cambridge University Press, Cam-